


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

The "AND" operator is unnecessary -- we include all search terms by default. [\[details\]](#)

Scholar [All articles](#) - [Recent articles](#) Results 1 - 10 of about 499 for (DC2 protein OR KIAA0404 protein) AND glioma

Protein kinase C involvement in apoptosis

M Lucas, V Sánchez-Margalet - General Pharmacology, 1995 - Elsevier

... dephosphorylation and activation of p34cdc2 does not ... I., Weiss MH and Law RE (1994)

Protein kinase C ... induce apoptosis in human malignant glioma cell lines ...

Cited by 93 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 4 versions

High Expression of Doublecortin and KIAA0369 Protein in Fetal Brain Suggests Their Specific Role in ... - [►amjpathol.org](#) (HTML)

M Mizuguchi, J Qin, M Yamada, K Ikeda, S Takashima - American Journal of Pathology, 1999 - ASIP

... cells, and U-251 and U-87 glioma cells, as ... Protein extraction, sodium dodecyl sulfate polyacrylamide electrophoresis, and ... 1200 for anti-DC1 and anti-DC2, 1:200 ...

Cited by 39 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 4 versions

Detection of protein kinase activity specifically activated at metaphase-anaphase transition -

[►jcb.org](#) (PDF)

M Sekimata - The Journal of Cell Biology, 1996 - Rockefeller Univ Press

... and KT34 immunoreactivities in U251 glioma cells. ... of protein utilizing the principle of protein-dye binding ... mitosis is mediated by p34cdc2 phosphorylation ...

Cited by 25 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 4 versions

Monocyte Chemoattractant Protein-1 (CCL2) in Inflammatory Disease and Adaptive Immunity: Therapeutic ...

C Daly, BJ Rollins - Microcirculation, 2003 - informaworld.com

... Given the fact that CCR2 is a G protein-coupled receptor, it is not ... these mice experience selective abnormalities in the trafficking of DC1 and DC2 cells (ie ...

Cited by 81 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 2 versions

... by a Recombinant Listeria monocytogenes Vaccine Targeting Tyrosinase Related Protein-2 and Real- ...

RM Prins, KW Bruhn, N Craft, JW Lin, CH Kim, SK ... - Neurosurgery, 2006 - neurosurgery-online.com

... numbers of nucleoprotein and tyrosinase related protein-2-specific ... a tumorigenic subline of the GL26 glioma deficient in ... cells, Dr. Ken Rock for the DC2.4 cells ...

Cited by 5 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 3 versions

... genes during permanent focal cerebral ischaemia: characterization of the protein kinase 9b5/MARKL1/ ...

A Schneider, R Laage, O von Ahsen, A Fischer, M ... - Journal of Neurochemistry, 2004 - pt.wkhealth.com

... org/cgi/data/298/5600/1912/ DC2/1 (Manning ... 9b5/MARK4 kinase, eg in competing with protein interactors. ... of the long variant MARK4L in different glioma cell lines ...

Cited by 21 - [Related articles](#) - [Web Search](#) - [BL Direct](#) - All 5 versions

Growth inhibition of glioma cells transfected with the human beta-interferon gene by liposomes ... - [►aacrjournals.org](#) (PDF)

M Mizuno - Cancer Research, 1990 - AACR

... $4.5 \pm 1.5 \times 10^5$ IU/cell (24.8 ± 3.8 ILJ/mg protein) was found ... E. Block of a glioma cell line in S by interferon ... Science (Washington DC), 2/9: 881-883, 1983 ...

[Cited by 61](#) - [Related articles](#) - [Web Search](#) - [All 3 versions](#)

Protein Transduction Domains Fused to Virus Receptors Improve Cellular Virus Uptake and Enhance ...

F Kuhnel, B Schulte, T Wirth, N Woller, S Schafers ... - Journal of Virology, 2004 - Am Soc Microbiol
... receptor to serve as molecular adapters between the adenoviral fiber knob protein and negatively ... DC2.4 cells were kindly provided by KL Rock (Worcester, Mass ...

[Cited by 9](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 6 versions](#)

Protein structure

SJ Winder, AE Knight, J Kendrick-Jones - Dystrophin: Gene, Protein and Cell Biology, 1997 - books.google.com
... full-length isoforms from skeletal muscle (Koenig et al., 1988) and glioma (Tinsley et al ... the first ~ 250 NH 2-terminal amino acids of each protein, giving an ...

[Cited by 10](#) - [Related articles](#) - [Web Search](#)

Immunotherapeutic Targeting of Shared Melanoma-Associated Antigens in a Murine Glioma Model - [aacrjournals.org](#) (HTML)

RM Prins, SK Odesa, LM Liao - Cancer Research, 2003 - AACR
... glioma cells, Dr. Ken Rock for DC2.4 cells ... intracranial; TRP-2, tyrosinase-related protein 2; MAA, melanoma-associated antigen; GAA, glioma-associated antigen ...

[Cited by 29](#) - [Related articles](#) - [Web Search](#) - [BL Direct](#) - [All 3 versions](#)

Key authors: [M Lucas](#) - [C Daly](#) - [B Nikolic](#) - [V Sánchez-Marg...](#) - [E Schneider](#)

Google 

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google